

ISSN : 0082 - 6340
E-ISSN : 2337 - 876X
Accredited : 30/E/KPT/2018



TREUBIA

*A JOURNAL ON ZOOLOGY
OF THE INDO-AUSTRALIAN ARCHIPELAGO*

Vol. 47, no. 1, pp. 1-52

June 2020



Published by

RESEARCH CENTER FOR BIOLOGY
INDONESIAN INSTITUTE OF SCIENCES
BOGOR, INDONESIA

ISSN : 0082 - 6340
E-ISSN : 2337 - 876X
Accredited : 30/E/KPT/2018



TREUBIA

*A JOURNAL ON ZOOLOGY
OF THE INDO-AUSTRALIAN ARCHIPELAGO*

Vol. 47, no. 1, pp. 1-52

June 2020



Published by

RESEARCH CENTER FOR BIOLOGY
INDONESIAN INSTITUTE OF SCIENCES
BOGOR, INDONESIA

ISSN : 0082 - 6340
E-ISSN : 2337 - 876X
Accredited : 30/E/KPT/2018

TREUBIA

A JOURNAL ON ZOOLOGY OF THE INDO-AUSTRALIAN ARCHIPELAGO
Vol. 47, no. 1, pp. 1–52, June 2020

Board of Editors:

Dr. Djunijanti Peggie, M.Sc. (Chief Editor)
Prof. Dr. Dewi Malia Prawiradilaga, M.Rur.Sc.
Dr. Daisy Wowor, M.Sc.
Dr. Kartika Dewi
Dr. Dhian Dwibadra
Dr. Conni Margaretha Sidabalok, M.App.Sc.

International Editors:

Dr. Paul Bates, M.A.	Harrison Institute Bowerwood House 15 Botolph's Road Sevenoaks, Kent, TN13 3AQ, UK
Dr. Thomas von Rintelen	Museum für Naturkunde Leibniz - Institut für Evolutions und Biodiversität Forschung an der Humboldt-University zu Berlin, Invalidenstraße 43, 10115 Berlin, Germany
Dr. Alan T. Hitch	University of California, Davis, CA 95616, USA

Reviewers:

Dr. R.I. Vane-Wright	Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation, University of Kent, Canterbury, CT2 7NR, UK; Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK
Dr. Chris J Müller	Australian Museum, 6 College Street, Sydney, NSW 2010, Australia
Dr. Willy Marthy Rombang	Wildlife Conservation Society-Indonesia Program, Jl. Malabar I no. 11, Bogor 16128, Indonesia
Dr. Ding Li Yong	BirdLife International (Asia), 354 Tanglin Road, Tanglin International Centre Singapore, 247672 Singapore
Prof. Dr. Dewi Malia Prawiradilaga, M.Rur.Sc.	Museum Zoologicum Bogoriense, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia
Muhammad Irham, M.Sc.	Museum Zoologicum Bogoriense, Research Center for Biology, Indonesian Institute of Sciences (LIPI), Indonesia

Managing Assistant:
Sri Wulan, S. Ikom.

Layout:
Liana Astuti

TREUBIA
RESEARCH CENTER FOR BIOLOGY - INDONESIAN INSTITUTE OF SCIENCES (LIPI)
Jl. Raya Jakarta-Bogor Km. 46, Cibinong-Bogor 16911, Indonesia
e-mail: treubia@gmail.com
<http://e-journal.biologi.lipi.go.id/index.php/treubia>

VOL. 47, NO. 1, JUNE 2020

CONTENT

David J. Lohman, Sarino, and Djunijanti Peggie Syntopic <i>Elymnias agondas aruana</i> female forms mimic different <i>Taenaris</i> model species (Papilionoidea: Nymphalidae: Satyrinae) on Aru, Indonesia	1–12
Tri Haryoko, Oscar Johnson, Matthew L. Brady, Subir B. Shakya, M. Irham, Yohanna, Rusdiyan P. Ritonga, Dewi M. Prawiradilaga, and Frederick H. Sheldon Recent ornithological expeditions to Siberut Island, Mt. Talamau and Rimbo Panti Nature Reserve, Sumatra, Indonesia	13–38
Elize Y. X. Ng, Arya Y. Yue, James A. Eaton, Chyi Yin Gwee, Bas van Balen, and Frank E. Rheindt Integrative taxonomy reveals cryptic robin lineage in the Greater Sunda Islands	39–52

TREUBIA

(A JOURNAL ON ZOOLOGY OF THE INDO-AUSTRALIAN ARCHIPELAGO)

ISSN : 0082 - 6340
E-ISSN : 2337 - 876X

Date of issue: 30 JUNE 2020

This abstract sheet may be reproduced without permission or charge

UDC: 595.78(594.73)

David J. Lohman

Syntopic *Elymnias agondas aruana* female forms mimic different *Taenaris* model species (Papilionoidea: Nymphalidae: Satyrinae) on Aru, Indonesia

TREUBIA, June 2020, Vol. 47, No. 1, pp. 1–12.

Wing patterns of female *Elymnias agondas* (Boisduval, 1832) butterflies are highly variable, presumably to mimic different *Taenaris* species throughout New Guinea and surrounding islands. Labels on most *E. agondas* museum specimens lack precise locality information, complicating efforts to match *E. agondas* female wing patterns with presumed *Taenaris* model species. This paucity of data also makes it impossible to determine where different forms occur and whether they are strictly allopatric. During fieldwork on the Aru Archipelago, we found two distinct forms of *E. agondas* females occurring syntopically. The “light form” resembles *T. catops*, while the “dark form” seems to mimic *T. myops* and *T. artemis*. We discuss the significance of this finding and illustrate species in the *Taenaris* mimicry ring encountered on Aru.

(David J. Lohman, Sarino, and Djunijanti Peggie)

Keywords: adaptation, Batesian mimicry, butterfly, mimicry ring, polymorphism

UDC: 598.2:910.4(594.4)

Tri Haryoko

Recent ornithological expeditions to Siberut Island, Mt. Talamau and Rimbo Panti Nature Reserve, Sumatra, Indonesia

TREUBIA, June 2020, Vol. 47, No. 1, pp. 13–38.

Siberut Island, Mt. Talamau, Rimbo Panti Nature Reserve, and intervening locations in West Sumatra Province were visited during two expeditions in 2018-2019 by ornithologists from the Museum Zoologicum Bogoriense-Indonesian Institute of Sciences (LIPI), Louisiana State University Museum of Natural Science, and Andalas University. The main objective of these expeditions was to obtain data and tissue-subsample rich museum specimens for morphological and genetic studies of phylogeny and population genetics of Southeast Asian birds aimed at understanding the causes of avian diversification in the region. We also observed, photographed, and audio-recorded numerous bird species during the expeditions and archived these data. In total, 285 species were identified, and specimen material was collected from 13 species and 26 subspecies not previously represented in tissue resource collections. Here, we provide complete lists of birds found at each location, highlight distributional discoveries, and note cases of potential taxonomic, ecological, and conservation interest.

(Tri Haryoko, Oscar Johnson, Matthew L. Brady, Subir B. Shakya, M. Irham, Yohanna, Rusdian P. Ritonga, Dewi M. Prawiradilaga, and Frederick H. Sheldon)

Keywords: birds, distribution, diversity, conservation, West Sumatra

UDC: 598.813.063(59)

Elize Y. X. Ng

Integrative taxonomy reveals cryptic robin lineage in the Greater Sunda Islands

TREUBIA, June 2020, Vol. 47, No. 1, pp. 39–52.

Southeast Asian avifauna is under threat from both habitat loss and illegal poaching, yet the region's rich biodiversity remains understudied. Here, we uncover cryptic species-level diversity in the Sunda Blue Robin (*Myiomela diana*), a songbird complex endemic to Javan (subspecies *diana*) and Sumatran (subspecies *sumatrana*) mountains. Taxonomic inquiry into these populations has previously been hampered by a lack of DNA material and the birds' general scarcity, especially *sumatrana* which is only known from few localities. We demonstrate fundamental bioacoustic differences in courtship song paired with important distinctions in plumage saturation and tail length that combine to suggest species-level treatment for the two taxa. Treated separately, both taxa are independently threatened by illegal poaching and habitat loss, and demand conservation action. Our study highlights a case of underestimated avifaunal diversity that is in urgent need of revision in the face of imminent threats to species survival.

(Elize Y. X. Ng, Arya Y. Yue, James A. Eaton, Chyi Yin Gwee, Bas van Balen, and Frank E. Rheindt)

Keywords: bioacoustics, bird trade, passerines, songbird crisis, taxonomic neglect

INTEGRATIVE TAXONOMY REVEALS CRYPTIC ROBIN LINEAGE IN THE GREATER SUNDA ISLANDS

Elize Y. X. Ng¹, Arya Y. Yue^{1,2}, James A. Eaton³, Chyi Yin Gwee¹, Bas van Balen⁴,
and Frank E. Rheindt^{*1}

¹Department of Biological Sciences, National University of Singapore, 16 Science Drive 4, 117558 Singapore

²Department of Biological Sciences, Bryn Mawr College, 101 N Merion Ave, Bryn Mawr, PA 19010, USA

³Casa Indah 1, Persiaran Surian, Petaling Jaya, 47810, Malaysia

⁴Basilornis Consults, Muntendampad 15, 6835 BE Arnhem, Netherlands

*Corresponding author: dbsrfe@nus.edu.sg

Received: 3 June 2020; Accepted: 26 June 2020

ABSTRACT

Southeast Asian avifauna is under threat from both habitat loss and illegal poaching, yet the region's rich biodiversity remains understudied. Here, we uncover cryptic species-level diversity in the Sunda Blue Robin (*Myiomela diana*), a songbird complex endemic to Javan (subspecies *diana*) and Sumatran (subspecies *sumatrana*) mountains. Taxonomic inquiry into these populations has previously been hampered by a lack of DNA material and the birds' general scarcity, especially *sumatrana* which is only known from few localities. We demonstrate fundamental bioacoustic differences in courtship song paired with important distinctions in plumage saturation and tail length that combine to suggest species-level treatment for the two taxa. Treated separately, both taxa are independently threatened by illegal poaching and habitat loss, and demand conservation action. Our study highlights a case of underestimated avifaunal diversity that is in urgent need of revision in the face of imminent threats to species survival.

Keywords: bioacoustics, bird trade, passerines, songbird crisis, taxonomic neglect

ABSTRAK

Fauna burung di Asia Tenggara mengalami ancaman yang disebabkan oleh kehilangan habitat dan perburuan liar, namun keanekaragaman tinggi di kawasan ini masih kurang dipelajari. Pada makalah ini kami mengungkapkan keragaman tersembunyi pada tingkat jenis burung berkecet biru tua (*Myiomela diana*), suatu kelompok burung berkicau yang pelik, endemik di pegunungan Jawa (anak jenis *diana*) dan Sumatra (anak jenis *sumatrana*). Sebelumnya penelitian taksonomi pada populasi burung tersebut terkendala oleh kekurangan material DNA dan kelangkaan burung, khususnya pada anak jenis *sumatrana* yang hanya diketahui dari sedikit lokasi. Kami menunjukkan perbedaan bioakustik dalam kicauan percumbuan dipasangkan dengan perbedaan penting pada warna bulu dan panjang ekor yang dikombinasikan untuk perlakuan tahap jenis pada kedua taksa. Diperlakukan sebagai jenis yang berbeda, kedua taksa ini terancam perburuan liar dan kehilangan habitat, dan memerlukan aksi konservasi. Studi ini menyoroti kasus keragaman fauna burung yang tidak terlalu diperhitungkan yang perlu segera direvisi dalam menghadapi ancaman nyata terhadap keberlangsungan jenis ini.

Kata kunci: bioakustik, perdagangan burung, passerines, krisis burung berkicau, pengabaian taksonomi

INTRODUCTION

Birds are the most well studied group of organisms. Yet despite this detailed taxonomic knowledge, there is ongoing turn-over in the classification of birds as a relatively high number of populations previously subsumed under more widespread species have recently gained recognition as independent species, especially in tropical regions of Asia (Collar, 2003; Barrowclough et al., 2016), indicating a potential underestimate of avian species diversity. Such erroneous assessments of avifaunal richness can impede conservation and

management actions that depend upon accurate taxonomic delimitations (Lohman et al., 2010; Rojas-Soto et al., 2010). Examples of such victims of unrecognized cryptic diversity include the Critically Endangered Javan Green Magpie (*Cissa thalassina*) that was once lumped with the non-threatened Bornean Green Magpie (*C. jefferyi*) (van Balen et al., 2013); the Critically Endangered Javan Leafbird (*Chloropsis cochinchinensis*) was previously considered a species with the widespread Blue-winged Leafbird (*Chloropsis moluccensis*) (BirdLife International, 2017); the Critically Endangered Javan Pied Starling (*Gracupica jalla*) was recognised as a species with its Least Concern northern sister taxon Asian Pied Starling (*Gracupica contra*) (BirdLife International, 2019).

In Southeast Asia, especially in the Greater Sundaic region, birds are threatened with habitat loss and illegal poaching (Symes et al., 2018). In particular, insular Southeast Asia has one of the highest rates of deforestation globally (Miettinen et al., 2011, FAO, 2015), a trend that is set to continue into the future as nations develop. To further exacerbate the critical situation, birds are trapped incessantly throughout the region for the illegal wildlife trade (Chng et al. 2015, 2016a, 2016b, 2018; Eaton et al. 2015, 2017a, 2017b; Krishnasamy & Stoner, 2016). There is a race to uncover cryptic species that remain beneath the conservation radar before they perish in the face of anthropogenic threats.

In this study, we uncover cryptic species-level diversity in the Sunda Blue Robin (*Myiomela diana*) complex from montane Java and Sumatra. The species consists of two currently recognized subspecies; *diana* from the western and central mountains of Java and *sumatrana* from the mountains of Sumatra (Fig. 1). Taxonomic inquiry into these two taxa has thus far been hampered by their general scarcity, especially *sumatrana* which is known from very few localities, and a general lack of DNA material. Both taxa have long been considered conspecific until Eaton et al. (2016a) upgraded them as separate species based on distinct vocalisations and plumage. Herein, we shed light on the taxonomic status of these two distinct taxa using an integrative approach relying on morphological specimen inspection and vocal analysis.

MATERIALS AND METHODS

We examined a subset of the series of *M. diana* specimens at the Natural History Museum at Tring (henceforth NHM Tring), Naturalis Museum (Leiden, Netherlands) (henceforth NM), and the Lee Kong Chian Natural History Museum (Singapore) (henceforth LKCNHM). In total we examined 30 specimens: 15 at NM comprising five male and five female *M. d. diana* and five male *M. d. sumatrana*; 12 at NHM Tring comprising six male and four female *M. d. diana* and one male and one female *M. d. sumatrana*; and 3 at LKCNHM comprising two male and one female *M. d. sumatrana* (Table S1). Five measurements were taken: (1) tarsus length, (2) length of upper mandible (as measured from the bill tip to the point where the upper mandible meets the forehead), (3) wing length, (4) tail length, and (5) total body length




Figure 1. Range map of the Sunda Blue Robin (*Myiomela diana*) complex. Illustration of Javan *diana* (bottom) and Sumatran *sumatrana* (top) depicting plumage differences in both males and females are modified from Eaton et al. (2016a).

(Table S1). Measurements (1) and (2) were taken with a calliper to the nearest tenth of a millimetre, while measurements (3), (4) and (5) were taken with a ruler to the nearest millimetre. Statistical analysis was not conducted for (1), (3) and (5) due to the large amount of missing measurements taken across examined specimens. Statistical differences between measurements were calculated using the one-sample Wilcoxon test as implemented in R3.4.3 (R Development Core Team, 2017). Plumage coloration of birds at NHM Tring was assessed against natural light. JAE took photos of NHM Tring specimens using a Sony DSC W90.

Sound recordings of courtship songs were obtained from online repository (Xeno-Canto) and comprise a total of 19 recordings which were included in our analysis (13 from *M. d. diana*, and six from *M. d. sumatrana*) (Table S2). We measured a total of nine vocal parameters: (1) number of elements per motif, (2) motif duration, (3) highest frequency, (4) lowest frequency, (5) bandwidth frequency, (6) peak frequency, (7) centre frequency (Charif et al., 2010), (8) frequency modulation in first half of motif, and (9) frequency modulation in second half of motif. Frequency modulations (parameters 8 and 9) were measured first by splitting the motif into four equal partitions, with parameter 8 reflecting the difference in peak frequency between the first and second partitions, and parameter 9 the difference between the third and the fourth. Vocal analyses were conducted and sonograms were generated using Raven Pro Version 1.5 (Bioacoustics Research Program, Cornell Laboratory of Ornithology, Ithaca, NY, USA). Default settings of Raven Pro were applied except window size, which was adjusted to 1000 after inspection of all sonograms to select the optimal level.

Principal Component Analysis (PCA) was conducted for vocal measurements in R using the ‘prcomp’ function. Additionally, we assessed the vocal diagnosability of variables using the criterion outlined by Isler et al. (1998), henceforth referred to as the Isler criterion. The Isler criterion is based on two conditions: (i) there must be no overlap between the ranges of measurements between the two taxa being compared, and (ii) the means \bar{x} and standard deviations (SD) of the taxon with the smaller set of measurements (a) and the taxon with the larger set of measurements (b) have to meet the following requirement: $\bar{x}_a + t_a SD_a \leq \bar{x}_b - t_b SD_b$, where t_i refers to the t-score at the 97.5th percentile of the t distribution for $n - 1$ degrees of freedom. Since there must be no overlap between the two sets of measurements, measurements of one taxon are uniformly higher than the other, and a one-tailed test with a significance level of 5% was used. The Isler criterion is substantially more discriminating than t-tests, Mann-Whitney U tests and tests including effect sizes because it uses the standard deviations of the sample points, not the standard deviation of the taxon mean, which is much smaller (Isler et al., 1998; Rheindt et al., 2011). In addition to the Isler criterion, a two-tailed t-test was also performed to assess the conservative nature of the Isler criterion.

RESULTS

Morphometrics

Both taxa were found to differ significantly in tail length. This difference was observed both in sex-specific comparisons (p-value = 0.003382 between males, and p-value = 0.03139 between females) and across all individuals (p-value = 0.005552) (Fig. 2, Table S1). There was no significant difference observed for bill length between the two taxa (p-value = 0.7011) (Fig. 2a). Based on total body length, *M. d. diana* is on average larger than *M. d. sumatrana*, both across and between sexes (Table S1).

Plumage

The male of *M. d. diana* has a glossy dark-blue plumage that can appear surprisingly pale when perched in the open in good light (JAE and FER, personal observation), strongly contrasting with the black primaries (Figs. 1 and S1). The small white feathers on the forecrown are usually hidden when the bird is feeding but often exposed when the bird is agitated or responding to playback in much the same way that the Javan Shortwing *Brachypteryx montana* can seemingly hide and expose its white supercilium on demand (JAE and FER, personal observation). Female *M. d. diana* shows bright chestnut upperparts and head with a small grey wedge on the lower throat, a rufous breast band and dark grey underparts across all specimens.

In *M. d. sumatrana*, the male is much darker blue, with little contrast with the black




Figure 2. Sex-specific box plots of morphometric measurements of (a) bill length (from bill tip to the point where the upper mandible meets the forehead); (b) tail length. Significant differences for the pairwise Wilcoxon Test are listed for $p \leq 0.05$ (*) and $p \leq 0.01$ (**). (c) Principal component (PC) plot based on nine vocal parameters for Sunda Blue Robins, with PC1 explaining 66.83% of total variance and PC2 explaining an additional 13.50% of total variance.

primaries (Fig. S1). The extent of white on the forecrown is similar to *diana*. Female *sumatrana* is also much darker in plumage compared to all female *diana*, lacking the bright chestnut tones (Fig. S1). Although the throat appears slightly paler than the breast and ear-coverts, it lacks the pronounced grey wedge of *diana*. The belly is a mucky, darker grey than female *M. d. diana* (Fig. S1).

Bioacoustics

The main courtship vocalizations of these two taxa are consistently and easily separable. *M. d. diana*'s song is a simple descending warble of 4-5 sweet melancholy glissading notes, usually with an ascending final note or – less commonly – a final flourish. In sharp contrast, *M. d. sumatrana*'s song is a much more heterogeneous, complex, tinkling motif – usually around twice as long as that of *diana* (Table S2) – consisting of a series of 5-8 high-pitched, rising-and-falling elements that invariably ascend into a fading pitch. This song is strongly reminiscent of the Bamboo Bush Warbler (=Yellow-bellied Warbler) *Abroscopus*

superciliaris.

Principal Component Analysis (PCA) of all bioacoustic measurements showed the two taxa to be distinctly different with no overlap in PCA space (Fig. 2c). It also agreed with our vocal characterizations in showing Sumatran songs to be more heterogeneous than Javan ones (i.e., filling out more bioacoustic space in Fig. 2c). Of the nine parameters, three – motif duration, highest and bandwidth frequency – did not overlap between the two taxa. However, due to the conservative nature of the Isler criterion, none of the parameters were deemed Isler-diagnosable (Table S2). In contrast, a two-tailed t-test revealed significant differences (p-values < 0.05, df = 18, t-value = 2.101) between the Javan and Sumatran taxa in 7 out of 9 parameters (except lowest and peak frequencies) (Table S2).

DISCUSSION

Many modern taxonomic treatments recognize the Sumatran and Javan taxa as subspecies of the Sunda Blue Robin *M. diana*. However, in light of the differences in plumage, vocalizations, and morphometrics here documented, we propose that the two subspecies should be recognized separately as Sumatran Blue Robin (*M. sumatrana*) and Javan Blue Robin (*M. diana*). The differences documented in plumage, vocalization, and morphometrics are testimony to the true differences between taxa, as both taxa have been showed to be statistically different despite being measured and/or recorded by various co-authors and/or recordists.

The subtle difference in plumage coloration is likely of reproductive significance, especially in the amount of male brightness, which may act as an honest signal of sexual selection (Hill, 2002; Dunn et al., 2015). More importantly, however, courtship vocalisations differ strongly between the two (Fig. 2c, Table S2). Vocalizations are an important factor in reproductive isolation in songbirds and have been used in documenting cryptic, overlooked species-level diversity in other Asian robins, thrushes and flycatchers (superfamily Muscicapoidae), e.g. *Brachypteryx* shortwings (Alström et al., 2018), *Sholicola* sholakilis (Robin et al., 2017), *Ficedula* flycatchers (Dong et al., 2015), and *Cyornis* jungle-flycatchers (Gwee et al., 2019).

In our taxonomic recommendation to separate *M. diana* and *M. sumatrana* into different species, we follow the yardstick approach (Mayr & Ashlock, 1991) under the umbrella of the multi-dimensional Biological Species Concept (Mayr, 1996). Although plumage differences detected are not substantial and may suggest mere subspecific differentiation, the important comparison is whether other Asian Muscicapoidae species pairs of relatively close phylogenetic affinity display plumage differences of the same order of magnitude or not. The same rationale can be applied to bioacoustic differences, which are arguably more important in reproductive isolation in these songbirds (Kroodsma & Miller, 1982; Kroodsma & Byers,

1991; Catchpole & Slater, 2003). The White-browed Shortwings of the *Brachypteryx montana* complex from mainland Asia have extensive similarities in plumage and have consequently been considered members of only one species until a combination of bioacoustic and molecular evidence led to their recent separation (Alström et al., 2018). In the same vein, vocal similarities arguably smaller than the ones observed between our robins have been documented among a number of *Cyornis* jungle-flycatchers (Gwee et al., 2019) that are known to be genetically deeply diverged (e.g. *Cyornis montanus* and an undescribed form from the Meratus mountains (Shakya et al., 2018; Eaton et al., 2016b). These cases illustrate that the vocal differences here documented between Javan and Sumatran Blue Robins are consistent with their treatment as independent species. Future studies can be made using genomic markers to confirm the distinctiveness between *M. diana* and *M. sumatrana*.

The recognition of Sumatran and Javan Blue Robins as independent species has important implications for their conservation. When cryptic species are lumped and treated as mere subspecies of each other because of taxonomic neglect, they are usually not granted the same conservation attention as compared to recognized species (Hazevoet, 1996). Population decline in such species often goes undetected as the overall population size of the combined umbrella species remains seemingly safe, leading to real extinction risk of species-level taxa (Hazevoet, 1996). Examples of such taxonomic neglect are abundant in the Indonesian Archipelago (e.g. Rheindt et al. 2017, 2020; Ng et al., 2016) and include Critically Endangered species such as the Javan Green Magpie (*Cissa thalassina*) (van Balen et al., 2013) and the Aceh Bulbul (*Pycnonotus bimaculatus*) (Eaton & Collar, 2015).

In the face of the songbird crisis that Southeast Asia is currently experiencing (Eaton et al., 2015), many species are threatened with extinction owing largely to the synergistic effects of illegal poaching for the bird trade and habitat loss (Symes et al., 2018). Currently, the Sunda Blue Robin is considered ‘Least Concern’ by the IUCN due to its large distribution on both Java and Sumatra (Birdlife International, 2018). With the present taxonomic revision, we propose that the Javan Blue Robin be given an IUCN classification of ‘Vulnerable’ under criterion A3 (IUCN, 2012) in which the population is projected or suspected to be reduced by at least 30% within the next 10 years. The species is restricted to the montane forests of west and central Java and is known to be targeted for the bird trade. A total of 19 *diana* individuals were found at three Jakarta markets during a single visit on one day (JAE, pers. obs., 4 July 2019). The Javan Blue Robin is only known from six localities across west and central Java (Fink et al., 2020), attesting to its rarity. Unsustainable poaching pressures will decimate populations if protection is not granted in the near future.

In a similar vein, we recommend that the Sumatran Blue Robin be given a classification of ‘Near Threatened’ based on the IUCN red list criteria: A3 (IUCN, 2012) – given the

increasing trapping pressures on Sumatran species as poachers seek to supply cage birds from elsewhere in the archipelago (Chng et al., 2015; Chng et al., 2018); B1a – despite the extensive reported range, the species has only been recorded at eight sites across Sumatra (Fink et al., 2020); B1bii – due to the ever increasing forest clearance in Sumatra that threatens the survival of the species (Miettinen et al., 2011). With habitat loss and illegal trade continuing to drive many Asian birds and other animals into extinction, it is increasingly important that taxonomic treatments reflect true species-level diversity as the survival of species depends upon it.

ACKNOWLEDGMENTS

We thank David Marquez, Mike Nelson, and Paulo Alves for contributing their recordings to Xeno-Canto. We additionally thank Rachel Oh and Yan Ru Choo for advice on statistical analysis. We are indebted to all museums and their staff including Naturalis Leiden, Kelvin Lim at the Lee Kong Chian Natural History Museum in Singapore, and Mark Adams at the Natural History Museum at Tring, for their kind assistance.

REFERENCES

- Alström, P., Rasmussen, P.C., Xia, C., Gelang, M., Liu, Y., Chen, G., Zhao, M., Hao, Y., Zhao, C., Zhao, C., Yao, C., Eaton, J.A., Hutchinson, R., Lei, F. & Olsson, U. 2018. Taxonomy of the White-browed Shortwing (*Brachypteryx montana*) complex on mainland Asia and Taiwan: an integrative approach supports recognition of three instead of one species. *Avian Research*, 9: 34.
- Alström, P., Saitoh, T., Williams, D., Nishiumi, I., Shigeta, Y., Ueda, K., Irestedt, M., Björklund, M. & Olsson, U. 2011. The Arctic warbler *Phylloscopus borealis* – three anciently separated cryptic species revealed. *Ibis*, 153: 395–410.
- van Balen, S.B., Eaton, J.A. & Rheindt, F.E. 2013. Biology, taxonomy and conservation status of the Short-tailed Green Magpie *Cissa [t.] thalassina* from Java. *Bird Conservation International*, 23: 91–109.
- Barrowclough, G.F., Cracraft, J., Klicka, J. & Zink, R.M. 2016. How many kinds of birds are there and why does it matter? *PLoS ONE*, 11: e0166307.
- BirdLife International. 2017. *Gracupica jalla* (amended version of 2017 assessment). *The IUCN Red List of Threatened Species 2017*: e.T103890801A118590020. <https://dx.doi.org/10.2305/IUCN.UK.2017-3.RLTS.T103890801A118590020.en>. 02 June 2020.
- BirdLife International 2018. *Myiomela diana*. ‘*The IUCN Red List of Threatened Species 2018*’: e.T22710114A132085301. <http://dx.doi.org/10.2305/IUCN.UK.2018-2.RLTS.T22710114A132085301.en>. 17 October 2019
- BirdLife International. 2019. *Chloropsis cochinchinensis*. *The IUCN Red List of Threatened Species 2019*: e.T103775551A156811213. <https://dx.doi.org/10.2305/IUCN.UK.2019-3.RLTS.T103775551A156811213.en>. 02 June 2020.
- Catchpole, C.K. & Slater, P.J. 2003. *Bird Song: Biological Themes and Variations*. Cambridge: Cambridge University Press: 348 pp.
- Charif, R.A., Waack, A.M. & Strickman, L.M. 2010. *Raven Pro 1.4 User’s Manual*. Cornell Lab of Ornithology, New York.

- Chng, S.C.L., Shepherd, C.R. & Eaton, J.A. 2018. In the market for extinction: birds for sale at selected outlets in Sumatra. *Traffic*, 30: 15–22.
- Chng, S.C.L. & Eaton, J.A. 2016a. Snapshot of an on-going trade: an inventory of birds for sale in Chatuchak weekend market, Bangkok, Thailand. *BirdingAsia*, 25: 24–49.
- Chng, S.C.L., Guciano, M. & Eaton, J.A. 2016b. In the market for extinction: Sukahaji, Bundung, Java, Indonesia. *BirdingAsia*, 26: 22–28.
- Chng, S.C.L., Eaton, J.A., Krishnasam, K., Shepherd, C.R. & Nijman, V. 2015. *In the Market for Extinction: An Inventory of Jakarta's Bird Markets*. TRAFFIC Southeast Asia, Selangor.
- Collar, N.J. 2003. How many bird species are there in Asia? *Oriental Bird Club Bulletin*, 38: 20–30.
- Dong, L., Wei, M., Alström, P., Huang, X., Olsson, U., Shigeta, Y., Zhang, Y. & Zheng, G. 2015. Taxonomy of the Narcissus Flycatcher *Ficedula narcissina* complex: an integrative approach using morphological, bioacoustic and multilocus DNA data. *Ibis*, 157: 312–325.
- Dunn, P.O., Armenta, J.K. & Whittingham, L.A. 2015. Natural and sexual selection act on different axes of variation in avian plumage color. *Science Advances*, 1: e1400155.
- Eaton, J.A. & Collar, N.J. 2015. The taxonomic status of *Pycnonotus bimaclatus snouckaerti*. *Forktail*, 31: 107–110.
- Eaton, J.A., Shepherd, C.R., Rheindt, F.E., Harris, J.B.C., van Balen, S.B., Wilcove, D.S. & Collar, N.J. 2015. Trade-driven extinctions and near-extinctions of avian taxa in Sundaic Indonesia. *Forktail*, 31: 1–12.
- Eaton, J.A., van Balen, B., Brickle, N.W. & Rheindt, F.E. 2016a. *Birds of Indonesian Archipelago: Greater Sundas and Wallacea*. 1st ed. Barcelona: Lynx Edicions: 496 pp.
- Eaton, J.A., Mitchell, S.L., Navario Gonzalez Bocos, C. & Rheindt, F.E. 2016b. A short survey of the Meratus Mountains, South Kalimantan province, Indonesia: two undescribed avian species discovered. *BirdingASIA*, 26: 107–113.
- Eaton, J.A., Leupen, B.T.C. & Krishnasamy, K. 2017a. *Songsters of Singapore: An overview of the bird species in Singapore pet shops*. TRAFFIC Southeast Asia, Selangor.
- Eaton, J.A., Nguyen, M.D.T., Willemsen, M., Lee, J. & Chng, S.C.L. 2017b. *Caged in the city: An inventory of birds for sale in Ha Noi and Ho Chi Minh City, Viet Nam*. TRAFFIC Southeast Asia, Selangor.
- Fink, D., Auer, T., Johnston, A., Strimas-Mackey, M., Robinson, O., Ligocki, S., Petersen, B., Wood, C., Davies, I., Sullivan, B., Iliff, M. & Kelling, S. 2020. *eBird Status and Trends, Data Version: 2018; Released: 2020*. Cornell Lab of Ornithology, Ithaca, New York.
- Food and Agriculture Organization of the United Nations (FAO). 2015 *Global Forest Resources Assessment 2015*. FAO, Rome.
- Gwee, C.Y., Eaton, J.A., Garg, K.M., Alström, P., van Balen, S., Hutchinson, R.O., Prawiradilaga, D.M., Le, H.M. & Rheindt, F.E. 2019. Cryptic diversity in *Cyornis* (Ave: Muscicapidae) jungle-flycatchers flagged by simple bioacoustic approaches. *Zoological Journal of the Linnean Society*, 186: 725–741.
- Hazevoet, C.J. 1996. Conservation and species lists: taxonomic neglect promotes the extinction of endemic birds, as exemplified by taxa from eastern Atlantic islands. *Bird Conservation International*, 6: 81–196.
- Hill, G.E. 2006. Female mate choice for ornamental coloration. In: Hill GE & McGraw KJ (ed) *Bird Coloration Vol II, Function and evolution*. pp 137–200. Harvard University Press: 528 pp.
- Isler, M.L., Isler, P.R. & Whitney, B.M. 1998. Use of vocalizations to establish species limits in antbirds (Passeriformes: Thamnophilidae). *Auk*, 115: 588–590.
- IUCN. 2012. IUCN Red List Categories and Criteria: Version 3.1. Second edition. IUCN, Gland, Switzerland and Cambridge, UK.
- Krishnasamy, K. & Stoner, S. 2016. *Trading faces: A rapid assessment on the use of Facebook to trade wildlife in Peninsular Malaysia*. TRAFFIC Southeast Asia, Selangor.

- Kroodsma, D.E. & Byers, B.E. 1991. The function(s) of bird song. *American Zoologist*, 31: 318–328.
- Kroodsma, D.E. & Miller, E.H. 1982. *Acoustic communication in birds*. Academic Press: 360 pp.
- Lohman, D.J., Ingram, K.K., Prawiradilaga, D.M., Winker, K., Sheldon, F.H., Moyle, R.G., Ng, P.K.L., Ong, P.S., Wang, L.K., Braile, T.M., Dwi Astuti & Meier, R. 2010. Cryptic genetic diversity in “widespread” Southeast Asian bird species suggests that Philippine avian endemism is gravely underestimated. *Biological Conservation*, 143: 1885–1890.
- Mayr, E. & Ashlock, P.D. 1991. *Principles of Systematic Zoology*, 2nd Edition. McGraw-Hill Inc: 475 pp.
- Mayr, E. 1996. What is a species, and what is not? *Philosophy of Science*, 63: 262–277.
- Miettinen, J., Shi, C. & Liew, S.C. 2011. Deforestation rates in insular Southeast Asia between 2000 and 2010. *Global Change Biology*, 17: 2261–2270.
- Ng, E.Y.X., Eaton, J.A., Verbelen, P., Hutchinson, R.O. & Rheindt, F.E. 2016. Using bioacoustic data to test species limits in an Indo-Pacific island radiation of *Macropygia* cuckoo doves. *Biological Journal of the Linnean Society*, 118: 786–812.
- Rheindt, F.E., Eaton, J.A. & Verbelen, F. 2011. Vocal trait evolution in a geographic leapfrog pattern: speciation in the Maroon-chinned Fruit Dove (*Ptilinopus subgularis*) complex from Wallacea. *Wilson’s Journal of Ornithology*, 123: 429–662.
- Rheindt, F.E., Christidis, L., Norman, J.A., Eaton, J.A., Sadanandan, K.R. & Schodde, R. 2017. Speciation in Indo-Pacific swiftlets (Aves: Apodidae): integrating molecular and phenotypic data for a new provisional taxonomy of the *Collocalia esculenta*. *Zootaxa*, 4250: 401–433.
- Rheindt, F.E., Prawiradilaga, M.D., Ashari, H., Suparno, Gwee, C.Y., Lee, G.W.X., Wu, M.Y. & Ng, N.S.R. 2020. A lost world in Wallacea: Description of a montane archipelagic avifauna. *Science*, 367: 167–170.
- R Core Team. 2017. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Robin, V.V., Vishnudas, C.K., Gupta, P., Rheindt, F.E., Hooper, D.M., Ramakrishnan, U. & Reddy, S. 2017. Two new genera of songbirds represent endemic radiations from the Shola Sky Islands of the Western Ghats, India. *BMC Evolutionary Biology*, 17: 31.
- Rojas-Soto, O.R., Navarro-Siguenza, A.G., de Los Monteros, A.E. 2010. Systematics and bird conservation policies: the importance of species limits. *Bird Conservation International*, 20: 176–185.
- Shakya, S.B., Haryoko, T., Burner, R.C., Prawiradilaga, D.M. & Sheldon, F.H. 2018. Preliminary assessment of community composition and phylogeographic relationships of the birds of the Meratus Mountains, south-east borneo, Indonesia. *Bulletin of the British Ornithology Club*, 138: 45–66.
- Symes, W.S., Edwards, D.P., Miettinen, J., Rheindt, F.E. & Carrasco, L.R. 2018. Combined impacts of deforestation and wildlife trade on tropical biodiversity are severely underestimated. *Nature Communications*, 9: 4052.

SUPPLEMENTARY MATERIAL

Table S1. Morphometric measurements for *M. d. diana* and *M. d. sumatrana*

Museum*	Taxon	Sex	Bill (mm)	Tarsus (mm)	Tail (mm)	Wing (mm)	Total length (mm)
NHM Tring	<i>sumatrana</i>	Male	14.5	-	58.5	83.1	139
NM	<i>sumatrana</i>	Male - Immature	16	26	59	-	125
NM	<i>sumatrana</i>	Male	17	25	no tail	-	-
NM	<i>sumatrana</i>	Male	16	25	63	-	137
NM	<i>sumatrana</i>	Male	17	26	59	-	134
NM	<i>sumatrana</i>	Male	15	26	58	-	160
LKCNHM	<i>sumatrana</i>	Male	-	26	61	81	146
LKCNHM	<i>sumatrana</i>	Male	19	26	66	82	153
<i>M. d. sumatrana</i> (Male) Mean			16.36	25.71	60.64	82.03	142.00
NHM Tring	<i>sumatrana</i>	Female	14.10	-	50.90	76.80	130.00
LKCNHM	<i>sumatrana</i>	Female	18.2	24.5	57	75	137
<i>M. d. sumatrana</i> (Female) Mean			16.15	24.50	53.95	75.90	133.50
<i>M. d. sumatrana</i> Mean			16.31	25.56	59.16	79.58	140.11
NHM Tring	<i>diana</i>	Male	15.1	-	70.1	84.9	160
NHM Tring	<i>diana</i>	Male	15	-	68.1	84.5	151
NHM Tring	<i>diana</i>	Male	15.5	-	69.7	87.8	162
NHM Tring	<i>diana</i>	Male	15.1	-	62.2	83	161
NHM Tring	<i>diana</i>	Male	14.8	-	67.2	85.2	160
NM	<i>diana</i>	Male	18	26	72	-	145
NM	<i>diana</i>	Male	18	28	74	-	150
NM	<i>diana</i>	Male	18	29	70	-	145
NM	<i>diana</i>	Male	16	26	69	-	140
NM	<i>diana</i>	Male	17	26	65	-	136
<i>M. d. diana</i> (Male) Mean			16.25	27.00	68.73	85.08	151.00
NM	<i>diana</i>	Female	16	26	63	-	140
NM	<i>diana</i>	Female	17	26	61	-	135
NM	<i>diana</i>	Female	17	25	68	-	145
NM	<i>diana</i>	Female	15	25	65	-	142
NM	<i>diana</i>	Female	17	28	61	-	132
NHM Tring	<i>diana</i>	Female	14.3	-	57.1	78.9	150
NHM Tring	<i>diana</i>	Female	14.6	-	65.9	80.3	155
NHM Tring	<i>diana</i>	Female	15.4	-	63.5	81.2	159
NHM Tring	<i>diana</i>	Female	13.9	-	62.5	80	140
NHM Tring	<i>diana</i>	Female	15	-	57.8	78.5	152
<i>M. d. diana</i> (Female) Mean			15.52	26.00	62.48	79.78	145.00
<i>M. d. diana</i> Mean			15.89	26.50	65.61	82.43	148.00

*Measurements made at NHM Tring (Natural History Museum at Tring, United Kingdom) were by JAE; NM (Naturalis Museum, Leiden, Netherlands) were by BvB; LKCNHM (Lee Kong Chian Natural History Museum, Singapore) were by CYG.

Table S2. Details of recordings used in this study. Mean for each recording measured is presented here

Taxon	Filename	Recordist	Location	No. of Motifs	Average no. of elements	Motif Duration (s)	Frequency (kHz)				Modulation		
							Highest	Lowest	Bandwidth	Peak	Centre	1st	2nd
<i>diانا</i>	Robin_Javan Blue_MN	Mike Nelson	Gunung Gede-Pangrango, Java	9	3.67	0.67	3859.88	2780.69	1079.19	3517.09	3517.09	0	0
	Robin_Javan Blue_MN 2	Mike Nelson	Gunung Gede-Pangrango, Java	14	4.00	0.64	4038.96	2571.13	1467.84	3374.54	3389.93	0	1
	Robin_Javan Blue_MN 3	Mike Nelson	Gunung Gede-Pangrango, Java	1	3.00	0.70	3760.90	2767.60	993.30	3186.90	3186.90	0	0
	Robin_Sunda Blue_Java	James Eaton	Gunung Gede-Pangrango, Java	22	4.09	0.65	4062.19	2744.40	1317.78	3717.40	3566.66	0	0
	Robin_Sunda Blue_Java_JE	James Eaton	Gunung Gede-Pangrango, Java	8	3.75	0.66	3889.84	2635.88	1253.96	3423.76	3439.90	1	1
	Cincidian5iii86_0714hrs_21Bxx4	Bas van Balen	Java	14	3.43	0.58	3976.43	2933.90	1042.53	3612.74	3545.76	0	0
	Cincidian17ii81_640_2B036	Bas van Balen	Java	7	3.71	0.55	4040.40	2798.89	1241.51	3582.60	3549.13	0	0
	20090822_GedePangrango_David Marquez	David Marquez Paulo	Gunung Gede-Pangrango, Java	5	4.80	0.71	3874.58	2680.90	1193.68	3367.78	3298.86	1	0
	Paulo Alves recordings 1	Alves Paulo	Java	3	3.33	0.56	3987.03	2798.53	1188.50	3287.37	3344.83	0	1
	Paulo Alves recordings 2	Alves Paulo	Java	15	3.73	0.63	3966.81	2760.84	1205.97	3471.15	3439.56	0	0
	Blue Robin_Java_Gede	-	Gunung Gede-Pangrango, Java	12	3.50	0.57	4026.14	2875.18	1150.96	3269.45	3280.21	0	1
	Myiomela diana_25v2019_735#98_Masigit	Bas van Balen	Mount Masigit, Java	12	3.67	0.57	4025.20	2790.92	1352.83	3562.50	3546.88	0	0
	Myiomela diana_4iii2019_622#87 Patuha (1)	Bas van Balen	Mount Patuha, Java	15	4.00	0.57	4022.55	2749.38	1422.83	3537.50	3425.00	0	0
	Min				3.00	0.55	3760.90	2571.13	993.30	3186.90	3186.90	0	0
	Max				4.80	0.71	4062.19	2933.90	1467.84	3717.40	3566.66	1	1
	SD				0.43	0.06	90.56	93.82	141.58	153.29	120.78	0.38	0.48
	mean				3.74	0.62	3963.92	2760.63	1223.91	3454.67	3425.44	0.15	0.31

Table S2. *continued*

Taxon	Filename	Recordist	Location	No. of Motifs	Average no. of elements	Motif Duration (s)	Frequency (kHz)				Modulation		
							Highest	Lowest	Bandwidth	Peak	Centre	1st	2nd
<i>sumatrana</i>	11_Mike Nelson_Sum_Kerinci	Mike Nelson	Gunung Kerinci, Sumatra	14	7.79	1.04	6638.58	2988.29	3650.29	3509.91	3512.97	1	1
	Robin_Sumatran Blue_MN	Mike Nelson	Gunung Kerinci, Sumatra	15	8.07	1.07	6198.68	3059.87	3138.81	3861.62	3738.15	1	1
	Robin_Sunda Blue_Kerinci	James Eaton	Gunung Kerinci, Sumatra	8	11.88	1.55	7087.13	2627.18	4459.95	4801.88	3886.73	1	1
	Robin_Sunda Blue_Sumatra	James Eaton	Gunung Kerinci, Sumatra	17	7.82	1.09	5696.61	2769.15	2927.46	3962.09	3886.11	1	1
	Cinc suma 201A188	Bas van Balen	Sumatra	5	4.00	1.06	4951.54	2998.86	1952.68	3562.50	3834.38	1	1
	Cinc suma203A339	Bas van Balen	Sumatra	5	5.00	0.76	4781.70	3010.96	1727.40	3646.88	3515.62	1	1
	Min			4.00	0.76	0.76	4781.70	2627.18	1727.40	3509.91	3512.97	1	1
	Max			11.88	1.55	1.55	7087.13	3059.87	4459.95	4801.88	3886.73	1	1
	SD			2.76	0.25	0.25	920.34	171.17	1028.77	479.24	174.90	0	0
	mean			7.43	1.10	1.10	5892.37	2909.05	2976.10	3890.81	3728.99	1	1
Isler Criterion			Criterion 1	Fail	Pass	Pass	Pass	Fail	Pass	Fail	Fail	Fail	Fail
			Criterion 2	-	Fail	Fail	Fail	-	Fail	-	-	-	-
Two-tailed t-test			p-value	0.02	0.01	0.01	0.00	0.09	0.01	0.08	0.01	0.00	0.00



Figure S1. Sunda Blue Robin males in (A) dorsal view and (B) ventral view. Sunda Blue Robin females in (C) dorsal and (D) ventral view. In A-D, *M. d. diana* (left) and *M. d. sumatrana* (right).

INSTRUCTIONS FOR AUTHORS

TREUBIA is a peer-reviewed, scientific zoological journal with focus on biosystematic aspects of terrestrial and aquatic fauna in the Indo-Australian region. TREUBIA is published twice a year and accepts manuscripts within the scope of the journal. It is accessible online at <http://e-journal.biologi.lipi.go.id/index.php/treubia>.

The missions of TREUBIA are to: (1) promote sciences and disseminate information in animal systematics and on the biodiversity of the region; (2) participate in the effort of educating public through good quality of scientific media and available professional researchers; (3) establish linkages among zoologists particularly in the field of systematics.

TREUBIA accepts manuscripts based on original research, taxonomic review or short communication. The manuscript should not be offered for prior or simultaneous publication elsewhere. It must be written in English and should use the American English spelling. Manuscripts should be prepared in Microsoft Word, using Times New Roman font 12, A4 paper size. Template is available through e-journal. An electronic file of the manuscript along with a formal cover letter – indicating the importance, stating its originality and its approval by all co-authors – should be submitted to the editors of TREUBIA through <http://e-journal.biologi.lipi.go.id/index.php/treubia> or through email address: treubia@gmail.com.

Concise writing is recommended. All numbers under 10 and any number forming the first word of a sentence must be spelled out, except in the Materials and Methods section of taxonomic papers. Year should be completely written. Names of genera and species should be in italic type. It is recommended to use metric measurements in abbreviation (for examples: kg, cm, ml). Please consult and refer to a recent issue of TREUBIA for an acceptable format. Please note that starting in 2018, we adopt Mendeley reference management application, with Harvard referencing style.

Manuscripts should be presented in the following order (with Conclusions and Appendices if necessary):

Title section. This includes the title of the paper (all capitalized), author's full name, author's institution and address (all with first letters capitalized), and e-mail address of the corresponding author. The title should be short, informative and without abbreviation.

Abstract. Except for short communications, articles should be accompanied by an abstract. The abstract consists of no more than 250 words in one paragraph which should clearly state the essence of the paper, with no references cited.

Keywords. Following the abstract, list up to 5 keywords, all typed in lowercase except a proper noun, separated by commas, presented in alphabetical order.

Introduction. The introduction must briefly justify the research and give the objectives. References related to the justification of the research should be cited in the introduction but extensive and elaborate discussion of relevant literature should be addressed in the Discussion section. References are to be cited in the text by the author's surname and year of publication. When citing multiple sources, place them in chronological order, for example: (Glaubrecht, 1999, 2006; Glaubrecht et al., 2009; Maaß & Glaubrecht, 2012). For two authors, both names should be cited. For three authors or more, only the first author is given followed by et al.

Materials and Methods. Provide a clear explanation of materials and methods used in the research. The place of specimen depository should be mentioned here.

Results. The results can be presented in the form of tables and figures when appropriate. The text should explain and elaborate the data presented. Captions of tables, figures, and plates should be inserted where you want them to be inserted. All line drawings, photographs and other figures should be submitted separately in JPEG format and the image size should be at least 1024 by 768 pixels.

Discussion. The discussion should interpret the results clearly and concisely, and should discuss the findings in relation with previous publications.

Acknowledgments. Acknowledgments of grants, assistance and other matters can be written in one paragraph.

References. List of references should be in alphabetical order by the first or sole author's surname. Journal references should include author's surname and initials, year of publication, title of the paper, full title of the journal (typed in *italic*), volume number and inclusive page numbers. Book references should include author's surname and initials, year of publication, title of the book (typed in *italic*) or/and title of the chapter and editor (if part of a book), publisher, city of publication, and page numbers.

For example:

Eaton, J.A., van Balen, B., Brickle, N.W. & Rheindt, F.E. 2016. *Birds of the Indonesian Archipelago: Greater Sundas and Wallacea*. 1st ed. Barcelona: Lynx Edicions.

LaSalle, J. & Schauff, M.E. 1994. Systematics of the tribe Euderomphalini (Hymenoptera: Eulophidae): parasitoids of whiteflies (Homoptera: Aleyrodidae). *Systematic Entomology*, 19: 235–258.

MacKinnon, J. & Phillips, K. 1993. *Field Guide to the Birds of Borneo, Sumatra, Java and Bali*. Oxford: Oxford University Press: 491 pp.

Natural History Museum 2013. Wallace100 - celebrating Alfred Russel Wallace's life and legacy. <http://www.nhm.ac.uk/nature-online/science-of-natural-history/wallace/index.html> 11 October 2013.

Higgins, P., Christidis, L., Ford, H. & Bonan, A. 2017. Honeyeaters (Meliphagidae). In: J. del Hoyo, A. Elliott, J. Sargatal, D.A. Christie & E. de Juana, eds. *Handbook of the Birds of the World Alive*. Barcelona: Lynx Edicions. <http://www.hbw.com>.

Upon receiving a manuscript, a Treubia editor will check the compliance with these instructions and will send the manuscript to two reviewers. Based on comments from the reviewers and the suitability of the manuscript, Treubia editors will decide the acceptance or rejection of the manuscript. The author will be notified of the decision and will receive the manuscript with reviewers' comments.

Following the process of reviewing and revising, a final proof will be sent to the first or sole author for correction and approval. Starting 2020, we will publish online only, without printed edition.

VOL. 47, NO. 1, JUNE 2020

CONTENT

David J. Lohman, Sarino, and Djunijanti Peggie Syntopic <i>Elymnias agondas aruana</i> female forms mimic different <i>Taenaris</i> model species (Papilionoidea: Nymphalidae: Satyrinae) on Aru, Indonesia	1–12
Tri Haryoko, Oscar Johnson, Matthew L. Brady, Subir B. Shakya, M. Irham, Yohanna, Rusdian P. Ritonga, Dewi M. Prawiradilaga, and Frederick H. Sheldon Recent ornithological expeditions to Siberut Island, Mt. Talamau and Rimbo Panti Nature Reserve, Sumatra, Indonesia	13–38
Elize Y. X. Ng, Arya Y. Yue, James A. Eaton, Chyi Yin Gwee, Bas van Balen, and Frank E. Rheindt Integrative taxonomy reveals cryptic robin lineage in the Greater Sunda Islands	39–52